

# COVERAGE OF MUAC SCREENING

Output indicator

---

## Indicator Phrasing

**English:** % of children aged 6 – 59 months screened for MUAC by the trained volunteers

**French:** % d'enfants âgés de 6 à 59 mois ayant fait l'objet d'un dépistage du PB par les volontaires formés

**Portuguese:** % de crianças com idades entre 6-59 meses triadas para o perímetro braquial (PB) pelos voluntários treinados

**Czech:** % dětí ve věku 6-59 let, kterým vyškolení dobrovolníci zkontrolovali obvod horní části paže

## What is its purpose?

The indicator measures the proportion of children screened in a given period by the trained volunteers (or other responsible personnel) for the prevalence of acute malnutrition (by measuring their mid-upper arm circumference, MUAC). The higher proportion of children is screened (and referred for treatment, if required), the higher proportion of undernourished children the treatment programs can identify and treat.

## How to Collect and Analyse the Required Data

Determine the indicator's value by using the following methodology:

- 1) Collect from the responsible volunteers in all the target villages the number of children screened during the last round of screening.
- 2) Divide the number by the total number of children aged 6 – 59 months living in the villages where the screening took place. Multiply the result by 100.
- 3) The result is the percentage of children screened for MUAC.

## Disaggregate by

Disaggregate the data by gender and specific vulnerable groups, such as minorities.

## Important Comments

### 1) **Use this indicator only if:**

- i) you have reliable and up-to-date statistics on the number of children aged 6-59.99 months living in the area where your screening system operates
- ii) the screening is done within a clearly defined period that does not overlap with other screening (i.e. there is a low risk of double counting)
- iii) you trust the data provided by health volunteers who conduct screening (for example, your supervisors randomly visit houses to check whether children were screened by volunteers)

Otherwise, you are likely to get unreliable data.

2) "Trained volunteer" means that the volunteer passed a knowledge and skills based performance test on measuring MUAC (i.e. is able to measure and record MUAC correctly).

3) This indicator relies on accurate age assessment. Since people often do not remember the exact dates of their children's birth, the data collectors should **always verify the child's age**. This can be done by reviewing the child's birth certificate, vaccination card or another document; however, since many caregivers do not have such documents (and since they can include mistakes), it is essential that your data collectors are able to **verify the child's age by using local events calendars**. Read FAO's Guidelines (see below) to learn how to prepare local events calendars and how to train data collectors in their correct use.

## Access Additional Guidance

- FAO (2008) [Guidelines for Estimating the Month and Year of Birth of Young Children](#)
- [Global Nutrition Cluster](#)