

AWARENESS OF FACILITIES WITH HIV TESTING

Output indicator

Indicator Phrasing

English: % of women aged 15 – 49 years aware of a place where they can get an HIV test

French: % de femmes âgées de 15 à 49 ans qui connaissent un endroit où elles peuvent effectuer un test VIH

Portuguese: % de mulheres com idades entre os 15 e os 49 anos que têm conhecimento de um local onde podem fazer o teste do VIH

Czech: % žen ve věku 15 – 49 let se znalostí místa kde mohou absolvovat HIV test

What is its purpose?

The indicator measures the proportion of women aware of where they can get an HIV test – an essential pre-condition for receiving effective treatment (if required) and preventing mother-to-child HIV transmission.

How to Collect and Analyse the Required Data

Collect the following data by conducting individual interviews with a [representative sample](#) of women aged 15 – 49 years:

RECOMMENDED SURVEY QUESTIONS (Q) AND POSSIBLE ANSWERS (A)

Q1: *Do you know of any place where you can get an HIV test?*

A1: yes / no

(ask only if the previous answer is YES)

Q2: *Where exactly is that place?*

A2: the respondent knows a specific place / the respondent does NOT know a specific place

Calculate the indicator's value by dividing the number of respondents who know a specific place by

the total number of interviewed respondents and multiplying the result by 100.

Disaggregate by

1) Always verify whether the respondent really **knows a specific place** and does not just say, for example, (somewhere) “in a district town” or (somewhere) “in a hospital”, without knowing a concrete place.

Important Comments

1) Always verify whether the respondent really **knows a specific place** and does not just say, for example, (somewhere) “in a district town” or (somewhere) “in a hospital”, without knowing a concrete place.